

CORRESPONDENCE

TIAPRIDE-INDUCED TORSADE DE POINTES

To the Editor:

Tiapride, an atypical neuroleptic agent, is a selective dopamine D2-receptor antagonist used for the treatment of agitation, aggressiveness, anxiety, and sleep disorders in elderly patients (1). We report a case of torsade de pointes, which has been associated with other neuroleptic agents (2–4), that occurred after the administration of tiapride to an elderly patient with agitation.

A 76-year-old man with atrial fibrillation was admitted for bronchitis and mild congestive heart failure. An electrocardiogram on admission showed atrial fibrillation at 47 beats per minute and a QTc interval of 440 msec. Treatment with intravenous furosemide and ceftriaxone was initiated. On day 3, oral tiapride (300 mg per day) was added for the treatment of agitation. On day 4, an electrocardiogram revealed a QTc interval of 600 msec. The 24-hour Holter recordings showed polymorphic ventricular tachycardia and QTc-interval prolongation. Thyroid hormone and serum electrolyte levels, including levels of potassium, magnesium, and calcium, were within normal ranges. After treatment with tiapride was discontinued, the QTc interval returned to normal (440 msec at a heart rate of 47 beats per minute) within 4 days. No other medication changes were made. One week later, results of a second 24-hour Holter monitor were normal.

In this patient, there was a strong temporal relation between tiapride administration and the development of QT prolongation and torsades de pointes. The patient had organic heart disease, which may have increased his propensity to develop torsade de pointes. No other cause, such as electrolyte imbalance, could be identified.

Acquired QT prolongation is caused primarily by drugs, and the list

of noncardiac drugs, including antibiotics, antimonial, and antifungal agents, prokinetic drugs, second-generation antihistamines, tricyclic antidepressants, and neuroleptic agents, continues to expand (5). Our case report suggests that tiapride administration may prolong the QT interval in patients with predisposing factors, such as atrial fibrillation and heart failure, and precipitate torsade de pointes. Tiapride is well tolerated at the doses recommended for elderly patients, but because of the potential risk of QTc-interval prolongation in patients with heart disease, medical supervision may be required.

Emilia Iglesias, MD

Ester Esteban

Sergio Zabala,

Antonio Gascon

Internal Medicine and Renal Unit

General Hospital Obispo Polanco

Teruel, Spain

1. Steele JW, Faulds D, Sorokin EM. Tiapride. A review of its pharmacodynamic and pharmacokinetic properties, and therapeutic potential in geriatric agitation. *Drugs Aging*. 1993;3:460–478.
2. Wilt JL, Minnema AM, Johnson RF, Rosenblum AM. Torsade de pointes associated with the use of intravenous haloperidol. *Ann Intern Med*. 1993;119:391–394.
3. Hunt N, Stern TA. The association between intravenous haloperidol and torsades de pointes, three cases and a literature review. *Psychosomatics*. 1995;36:541–549.
4. Sharma ND, Rosman HS, Padhi ID, Tisdale JE. Torsades de pointes associated with intravenous haloperidol in critically ill patients. *Am J Cardiol*. 1998;81:238–240.
5. Yap YG, Camm J. Risk of torsades de pointes with non-cardiac drugs. Doctors need to be aware that many drugs can cause QT prolongation. *BMJ*. 2000;320:1158–1159.

HEARING LOSS DUE TO ACUTE LEUKEMIA

To the Editor:

Ear involvement has rarely been described in patients with acute or chronic leukemia (1). Otologic symptoms are usually related to bleeding, tumoral infiltration, or infection. We

report 2 patients in whom hearing loss led to a diagnosis of acute leukemia.

A 31-year-old man was admitted with a 1-month history of severe left external otitis. Laboratory examination revealed a leukocyte count of $5.3 \times 10^9/L$ with 28% blasts, $37 \times 10^9/L$ platelets, and a hemoglobin level of 91 g/L. Examination of the bone marrow established a diagnosis of acute myeloid leukemia (type 2 FAB). The patient underwent chemotherapy, followed by autologous bone marrow transplantation. A complete hematologic remission was obtained and the otitis resolved. Eight months later, left external otitis recurred, and a blood count confirmed a relapse of leukemia. Salvage chemotherapy led to partial remission and resolution of the otitis. One month later, the patient presented with external otitis, hearing loss, and uncompleted left facial palsy. Computed tomographic (CT) scans and magnetic resonance imaging (MR) images of the left temporal bone showed diffuse inflammatory content in the tympanic cavity and adjacent air cells, without destruction of the ossicles and labyrinth (Figure 1). Surgical exploration revealed a left middle-ear chloroma. Palliative external irradiation (30 Gy) achieved local control, and oral chemotherapy was administered, with supportive care. The patient died 3 months later of disseminated invasive aspergillosis.

A 55-year-old man experienced sudden left sensorineural hearing loss followed by progressive right conductive hearing loss during the next 2 months. He was diagnosed with otitis media, but therapy with oral antibiotics was ineffective. He was then admitted due to the onset of trigeminal neuralgia. Examination revealed bilateral otitis media with a thickened ear drum. The skin of the external auditory canal was thickened as well. Right myringotomy showed a bloody effusion of which the culture was sterile. A CT scan showed that the mid-