

Table 2 Propensity Score Analyses of Relative Risk of In-Hospital Mortality in the Filter Group Versus the No-filter Group

	Filter Group		No-filter Group		P	Risk Ratio (95% CI)	Risk Difference (95% CI)	NNT (95% CI)
	No. of Deaths / Total No.	%	No. of Deaths / Total No.	%				
Unadjusted (n = 13,125)	97/3948	2.5%	522/9177	5.7%	<.001	0.43 (0.35-0.53)	-3.2% (-3.9% to -2.6%)	31 (26-39)
1:1 Propensity score-matched (n = 6948)	91/3474	2.6%	164/3474	4.7%	<.001	0.55 (0.43-0.71)	-2.1% (-3.0% to -1.2%)	48 (34-82)
Inverse probability of treatment weighting (n = 26,230)	354/13,106	2.7%	704/13,124	5.4%	<.001	0.50 (0.44-0.57)	-2.7% (-3.1% to -2.2%)	38 (32-46)

CI = confidence interval; NNT = number needed to treat.

Cohen SM, Kim J, Roy N, et al. Delayed Otolaryngology Referral for Voice Disorders Increases Health Care Costs. *Am J Med.* 2015; 128:426.e11-426.e18.



Three values listed in **Table 3** of this article were erroneous. The corrected values appear in bold type in this revised table.

Table 3 Mean and Median Annual Outpatient Costs per Person Involved in the Evaluation and Management of Laryngeal/Voice Disordered Patients by Time from First Primary Care Outpatient Visit to First Outpatient Otolaryngology Visit

	≤1 Month	>1 Month and ≤3 Months	> 3 Months
Preotolaryngology costs*			
Mean (SD)	224.20 (489.47)	274.94 (611.00)	333.55 (846.84)
Median	79.37	92.63	84.27
25 th percentile	54.96	59.95	57.57
75 th percentile	181.53	242.76	204.88
% Medication costs	69.7%	68.9%	71.6%
% Outpatient visit costs	30.3%	31.1%	28.4%
% Procedure costs	0%	0%	0%
Postotolaryngology costs*			
Mean (SD)	1598.11 (3714.62)	1950.97 (4550.59)	2379.64 (5183.15)
Median	729.63	842.84	1015.90
25 th percentile	378.50	417.01	424.14
75 th percentile	1592.92	1882.46	2358.04
% Medication costs	19.7%	22.7%	51.8%
% Outpatient visit costs	26.4%	20.4%	14.9%
% Procedure costs	53.9%	46.0%	33.3%
Total costs*			
Mean (SD)	1822.31 (3821.93)	2225.92 (4725.28)	2713.19 (5544.95)
Median	906.72	1037.58	1263.16
25 th percentile	470.57	537.00	611.67
75 th percentile	1918.00	2237.79	2684.44
% Medication costs	25.8%	38.01%	54.3%
% Outpatient visit costs	26.9%	21.7%	16.6%
% Procedure costs	47.3%	40.3%	29.2%

*Preotolaryngology costs = costs incurred from first primary care outpatient visit up to but not including the first otolaryngology outpatient visit. Postotolaryngology costs = costs incurred from first otolaryngology outpatient visit through 6 months after this visit. Total costs = preotolaryngology + postotolaryngology costs.

Christianson L, Matthews RV, Rahmitoola SH. Early Repolarization Masquerading as STEMI. *Am J Med.* 2014;127:e1-e2.



The last author's last name should be Rahimtoola SH.